

Tasectan[®]

Intended use

Tasectan is a medical device used to restore the physiological function of the intestinal walls. It is specifically formulated to control and reduce the symptoms associated with diarrhoea resulting from various causes, such as abdominal tension and frequent defecation. It is effective within 12 hours.

The Gelatin Tannate of which the product is composed is not altered in the stomach and acts by forming a film which protects the intestinal mucosa, reducing the frequency and duration of diarrhoea episodes. The medical device is available in capsule form for adults and as a powder recommended for paediatric use.

Nature and contents of the container

- Twenty 250 mg sachets (paediatric use).
- Blister pack containing fifteen 500 mg capsules.

Composition

Capsules: Gelatin Tannate, Corn Starch, Magnesium Stearate

Powder for paediatric use: Gelatin Tannate

Instructions for use

Capsules: swallow the capsule with water.

Powder: it can be added to food or mixed with a small quantity of milk, fruit juices or water. Mix well and give immediately.

Dosage

Adults: 1-2 capsules every 4-6 hours until the symptoms disappear.

Children under 3 years: 1 sachet every 6 hours until the symptoms disappear.

Children aged 3 to 14 years: 1 or 2 sachets every 6 hours until the symptoms disappear.

Warnings

- The consultation of a healthcare professional before using the product is not necessary. However such consultation is advisable in case of severe or persistent symptoms or when there are doubts on the diagnosis, mainly in children below 3 years and elderly people.
- In case of diarrhoea in children below 3 years, mainly when other symptoms are present, we recommend to consult your healthcare professional
- It's recommended abundant liquid intake and the usual dietary measures used in the management of diarrhoea (oral rehydration).
- Even though no side effects are known, it is recommended that the product is not used during pregnancy or in the early months of breastfeeding. In any case, always consult your doctor beforehand.
- Do not use the product after the expiry date printed on the package.
- Do not use the product if the blister or sachets is opened or damaged
- Store at room temperature away from sources of heat. Do not freeze.
Keep the product out of the reach of children.

Contraindications and undesired effects

Tasectan should not be used in patients with known hypersensitivity to gelatin tannate or any other ingredient of the product.



NOVINTETHICAL PHARMA SA
Via Pian Scairolo, 11
6915 Lugano - Switzerland

CE
0373

Rev:04-b dated 05.02.2013