

# Tasectan Duo<sup>®</sup>

## Intended use

The medical device is used to restore the physiological function of the intestinal wall as well as to prevent and alleviate dysbacteriosis of different nature and origin.

It is indicated for the prevention and treatment of digestive disorders as diarrhoea and other related symptoms such as bloating and abdominal tension, as a result of infectious gastroenteritis, antibiotic treatments and chemotherapy among others, being effective within the first 12 hours. The medical device contributes to restore the optimal gut conditions to promote growth of the endogenous intestinal flora thus contributing to improve the natural defences of the organism.

## Instruction for use

Powder for pediatric use: it has to be added to food or mixed with a small amount of milk, fruit juices or water. Mix well and give immediately.

Tablets for adult use: swallow the tablets with water.

## Contents of the container

Powder for pediatric use: 12 sachets containing gelatin tannate and tyndalized lactic bacteria.

Tablets for adult use: 12 tablets containing 500 mg of Gelatin Tannate and tyndalized lactic bacteria.

## Composition

Powder: Gelatin Tannate, Tyndalized Lactic Bacteria

Tablets: Gelatin Tannate, Tyndalized lactic bacteria, Calcium phosphate, Microcrystalline precipitated silica, Manesium stearate, Croscarmellose sodium, Polyvinylpyrrolidone, Hydroxypropylmethyl cellulose, Titanium dioxide, Microcrystalline cellulose, Stearic acid, Glycerol, Indigo carmine E 132 and Indigo carmine aluminium lake, Purified water, Vegetable glycerin.

Sources: Lactobacillus acidophilus, Lactobacillus plantarum, Lactobacillus casei, Lactobacillus rhamnosus, Bifidobacterium bifidum, Streptococcus thermophilus

## Dosage

For children under 3 years old: 1 sachet every 6 hours until the symptoms disappear

For children over 3 years old: 1 or 2 sachets every 6 hours until the symptoms disappear

1-2 tablets every 4-6 hours until the symptoms disappear.

## Warnings

Please note the following warnings

- The consultation of a healthcare professional before using the product is not necessary. However such consultation is advisable in case of severe or persistent symptoms or when there are doubts on the diagnosis, mainly in children below 3 years and elderly people.
- In case of diarrhoea in children below 3 years, mainly when other symptoms are present, we recommend to consult your healthcare professional
- It's recommended abundant liquid intake and the usual dietary measures used in the management of diarrhoea (oral rehydration).
- Even though no side effects are known, it is recommended that the product is not used during early months of pregnancy or during breastfeeding. In any case, always consult your doctor beforehand.
- Do not use the product after the expiry date printed on the package.
- Do not use the product if the blister or sachets are opened or damaged
- Store at room temperature away from sources of heat. Do not freeze.
- Keep the product out of the reach of children.

## Contraindications and undesired effects

Do not use the medical device in case of allergy to gelatin tannate or any ingredient of the product.



NOVINTETHICAL PHARMA SA  
Via Pian Scairolo, 11  
6915 Lugano - Switzerland



Rev 06: 12.12.2014