

LEAFLET

Tasectan® Duo

Intended use

Tasectan Duo is used to restore the physiological function of the intestinal wall as well as to prevent and alleviate dysbacteriosis of different nature and origin.

It is indicated for the prevention and treatment of digestive disorders as diarrhoea and other related symptoms such as bloating and abdominal tension, as a result of infectious gastroenteritis, antibiotic treatments and chemotherapy among others, being effective within the first 12 hours. The medical device contributes to restore the optimal gut conditions to promote growth of the endogenous intestinal flora thus contributing to improve the natural defences of the organism.

Presentations

Twelve sachets containing powder for paediatric use.

Blister pack containing twelve tablets for adult use.

Composition

Powder for paediatric use: 250 mg Gelatin Tannate (Porcine origin), Tyndalized Lactic bacteria.

Tablets: 500 mg Gelatin Tannate (Porcine origin), Tyndalized Lactic bacteria, Calcium Phosphate, Amorphous Precipitated Silica, Magnesium Stearate (Vegetable origin), Croscarmellose Sodium, Polyvinylpyrrolidone, Hydroxypropylmethyl Cellulose, Titanium Dioxide, Microcrystalline Cellulose, Stearic Acid, Indigo Carmine E 132 and Indigo Carmine Aluminium Lake, Vegetable Glycerin.

Sources: *Lactobacillus acidophilus*, *Lactobacillus plantarum*, *Lactobacillus casei*, *Lactobacillus rhamnosus*, *Bifidobacterium bifidum*, *Streptococcus thermophilus*.

Instructions for use

Powder: the content of one sachet must be mixed with a small quantity of a not hot drink such as milk, fruit juices or water and administered immediately. It can also be added to yoghurt or any other food. Do not administer the product directly without mixing it.

Tablets: Swallow the tablet with liquids.

Dosage

Children under 3 years: 1 sachet (paediatric use) every 6 hours until symptoms disappear.

Children aged 3 to 14 years: 1 or 2 sachets (paediatric use) every 6 hours until symptoms disappear.

Adults and children older than 14 years: 1-2 tablets every 4-6 hours until symptoms disappear.

Warnings

- Consultation with a healthcare professional before using the product is not necessary. However, such consultation is advisable in case of severe or persistent symptoms or when there are doubts on the diagnosis.
- In the case of diarrhoea in children below 3 years and elderly people, consultation with a healthcare professional is recommended, especially when other symptoms are present.
- This medical device is not a pharmacological treatment. If such treatment is recommended by a healthcare professional, the device can be administered concomitantly.
- An abundant intake of liquid and dietary measures accepted in the management of diarrhoea is recommended.
- Even though no side effects are known, it is recommended that the product is not used during pregnancy or breastfeeding, unless otherwise indicated by a healthcare professional.
- Do not use the product after the expiry date printed on the package.
- Do not use the product if the blister or sachets are opened or damaged
- This medical device does not require special storage conditions. Do not freeze.
- Keep product out of reach of children.

Contraindications

Tasectan Duo must not be used in patients with known hypersensitivity to Gelatin Tannate or any other ingredient of the product.



NOVENTURE, S.L
Avenida Diagonal, 549, 5a planta
08029 Barcelona
España

CE
0373

Rev. 04: 01.09.2019