

LEAFLET

Utipro® Plus

Intended use

Utipro Plus is intended for the control and prevention of urinary tract infections (cystitis) produced by pathogens such as *E.coli* and other gram (-) bacteria normally involved in the etiology of urological infections.

The compound, based on Xyloglucan (Hemicellulose) and Gelatin, acts mechanically in the intestinal lumen, avoiding the contact of the pathogens with the intestinal mucosa, which is the first step for their proliferation and subsequent passage to the urinary tract.

Utipro Plus is to be taken orally, from the appearance of the first symptoms of urinary tract disorders, with the aim of reducing the proliferation of pathogens that may degenerate into infections. In case of infection recurrences, Utipro Plus is also intended for the prevention of new episodes, in which the proliferation of pathogen cause repeated urinary tract infection.

Presentation

Blister pack containing 15 or 30 capsules.

Composition

Each capsule contains: 100 mg of Xyloglucan, 50 mg of Gelatin (porcine origin), 100 mg of Propolis, 100 mg of *Hibiscus sabdariffa*, Silicon Dioxide, Magnesium Stearate (vegetable origin) and Corn Starch.

Instructions for use

Swallow the capsule with liquids.

Dosage

In patients who develop the first urinary discomfort symptoms, Utipro Plus must be taken orally as 2 capsules per day (1 capsule every 12 hours) for five days.

For the prevention of recurrence, one capsule must be taken per day for at least 15 consecutive days per month. If required, the treatment can be repeated

Warnings

- The diagnosis of a urinary tract infection (UTI) should be made by a physician. If symptoms persist for more than 5 days, consult your physician.
- An acute UTI may require treatment with antibiotics. This product does not have bactericidal or bacteriostatic properties and, therefore, cannot substitute antibiotic treatment. However, Utipro Plus can be administered concomitantly.
- It is recommended that the product is not used during pregnancy or breastfeeding, unless otherwise indicated by a healthcare professional.
- Do not use the product after the expiry date printed on the package.
- Do not use the product if the blister is opened or damaged.
- Do not store above 25°C. Do not freeze
- Keep the product out of reach of children.

Contraindications

Utipro Plus must not be used in patients with known hypersensitivity to Xyloglucan, Gelatin or any other ingredient of the product.



NOVENTURE, S.L
Avenida Diagonal, 549, 5ª planta
08029 Barcelona
España

CE
0373

Rev.04:01.09.2019

|

LEAFLET

Utipro® Plus AF

Intended use

Utipro Plus AF is intended for the control and prevention of urinary tract infections produced by pathogens such as *E.coli* and other gram (-) bacteria normally involved in the etiology of urological infections.

The compound, based on Xyloglucan (Hemicellulose) and Gelose, acts mechanically in the intestinal lumen, avoiding the contact of the pathogens with the intestinal mucosa, which is the first step for their proliferation and subsequent passage to the urinary tract.

Utipro Plus AF is to be taken orally, from the appearance of the first symptoms of urinary tract disorders, with the aim of reducing the proliferation of pathogens that may degenerate into infections. In case of infection recurrences, Utipro Plus AF is also intended for the prevention of new episodes, in which the proliferation of pathogen cause repeated urinary tract infection.

Presentation

Blister pack containing 15 capsules.

Composition

Xyloglucan, Gelose, Propolis, *Hibiscus sabdariffa*, Silicon Dioxide, Magnesium Stearate (vegetable origin) and Corn Starch.

Instructions for use

Swallow the capsule with liquids.

Dosage

In patients who develop the first urinary discomfort symptoms, Utipro Plus AF must be taken orally as 2 capsules per day (1 capsule every 12 hours) for five days.

For the prevention of recurrence, one capsule must be taken per day for at least 15 consecutive days per month. If required, the treatment can be repeated.

Warnings

- The diagnosis of a urinary tract infection (UTI) should be made by a physician.
- An acute UTI may require treatment with antibiotics. This product does not have bactericidal or bacteriostatic properties and, therefore, cannot substitute antibiotic treatment. However, Utipro Plus AF can be administered concomitantly.
- Even though no side effects are known, it is recommended that the product is not used during pregnancy or breastfeeding, unless otherwise indicated by a healthcare professional.
- Do not use the product after the expiry date printed on the package.
- Do not use the product if the blister is opened or damaged.
- Do not store above 25°C. Do not freeze.
- Keep the product out of reach of children.

Contraindications

Utipro Plus AF must not be used in patients with known hypersensitivity to Xyloglucan or any other ingredient of the product.



NOVENTURE, S.L
Avenida Diagonal, 549, 5ª planta
08029 Barcelona
España

CE
0476

