

# **LEAFLET**

## **Xilaplus®**

### **Intended use**

Xilaplus is intended to restore the physiological functions of the intestinal walls, specifically formulated for the control and reduction of the symptoms related to diarrheal events of different aetiologies, such as abdominal tension and frequent emission of faeces.

The combination of Xyloglucan and Gelatin of which the product is composed acts by forming a film which protects the intestinal mucosa, reducing the frequency and duration of diarrhoea episodes.

### **Presentations**

Eight sachets containing powder for paediatric use.

Blister pack containing eight capsules for adult use.

### **Composition**

Powder: Combination of Xyloglucan and Gelatin (porcine origin) Capsules: Combination of Xyloglucan and Gelatin (porcine origin), Corn Starch, Magnesium Stearate (Vegetable origin).

### **Instructions for use**

Powder: Dissolve the contents of one sachet in half a glass of water. It can also be added to a small quantity of fruit juice. Mix well and administer immediately.

Capsules: Swallow the capsule with liquids.

### **Dosage**

Children under 3 years: 1 sachet every 8 hours.

Children aged 3 to 14 years: 1-2 sachets every 6-8 hours.

Adults and children older than 14 years: 1-2 capsules every 6-8 hours.

### **Warnings**

- Consultation with a healthcare professional before using the product is not necessary. However, such consultation is advisable in case of severe or persistent symptoms or when there are doubts on the diagnosis.
- In the case of diarrhoea in children below 3 years and elderly people, consultation with a healthcare professional is recommended, especially when other symptoms are present.
- This medical device is not a pharmacological treatment. If such treatment is recommended by a healthcare professional, the device can be administered concomitantly.
- An abundant intake of liquid and dietary measures accepted in the management of diarrhoea is recommended.
- Even though no side effects are known, it is recommended that the product is not used during pregnancy or breastfeeding, unless otherwise indicated by a healthcare professional.
- Do not use the product after the expiry date printed on the package.
- Do not use the product if the blister or sachets are opened or damaged.
- This medical device does not require special storage conditions. Do not freeze.
- Keep product out of reach of children.

### **Contraindications**

Xilaplus must not be used in patients with known hypersensitivity to Xyloglucan, Gelatin or any other ingredient of the product.



NOVENTURE, S.L  
Avenida Diagonal, 549, 5ª planta  
08029 Barcelona  
España

CE  
0373

Rev.05: 01.09.2019

# **LEAFLET**

## **Xilaplus® AF**

### **Intended use**

The medical device is intended to restore the physiological functions of the intestinal walls, specifically formulated for the control and reduction of the symptoms related to diarrheal events of different aetiologies, such as abdominal tension and frequent emission of faeces. The medical device acts by forming a film which protects the intestinal mucosa, reducing the frequency and duration of diarrhoea episodes.

### **Presentations**

Eight sachets containing powder for paediatric use with flavor.

Eight or Twelve sachets containing powder for paediatric use without flavour.

Blister pack containing eight capsules for adult use.

Blister pack containing eight tablets for adult use.

### **Composition**

Powder for pediatric use with flavour: Xyloglucan, Gelose, Maltodextrins, Citric acid, Sodium Chloride, Sucralose and Strawberry and Vanilla Flavours.

Powder for pediatric use without flavour: Xyloglucan, Gelose, Silicon Dioxide.

Capsules: Xyloglucan, Gelose, Magnesium Stearate (vegetable origin), Silicon Dioxide

Tablets: Xyloglucan, Gelose, Cellulose, Magnesium Stearate (vegetable origin), Silicon Dioxide.

### **Instructions for use**

Powder for pediatric use with flavour: Dissolve the content of one sachet in a half glass of water. It can also be added to fruit juices. Mix well and give immediately.

Powder for pediatric use without flavour: mix the content of one sachet with a small amount of water and give immediately. Do not administer the product directly without mixing it.

Capsules: swallow the capsule with liquids.

Tablets: swallow the tablet with liquids.

### **Dosage**

Children under 3 years: 1 sachet every 8 hours.

Children above 3 years: 1-2 sachets every 6-8 hours.

Adults: 1-2 tablets or capsules every 6-8 hours.

### **Warnings**

- Consultation with a healthcare professional before using the product is not necessary. However, such consultation is advisable in case of severe or persistent symptoms or when there are doubts on the diagnosis.
- In the case of diarrhoea in children below 3 years and elderly people, consultation with a healthcare professional is recommended, especially when other symptoms are present.
- This medical device is not a pharmacological treatment. If such treatment is recommended by a Healthcare professional, the device can be administered concomitantly with it.
- An abundant intake of liquid and dietary measures accepted in the management of diarrhoea is recommended.
- Even though no side effects are known, it is recommended that the product is not used during pregnancy or breastfeeding, unless otherwise indicated by a healthcare professional.

- Do not use the product after the expiry date printed on the package.
- Do not use the product if the blister or sachets are opened or damaged.
- This medical device does not require any special storage conditions. Do not freeze.
- Keep product out of reach of children.

### **Contraindications**

The medical device should not be used in patients with known hypersensitivity to any ingredient of the product listed in composition.



**NOVENTURE, S.L**

Avenida Diagonal, 549, 5ª planta  
08029 Barcelona  
España

CE

0476

Rev.06:01.09.2019